

Topical Steroid Misuse: An Unrecognized Cause Of Iatrogenic Cushing Syndrome In Infancy

Verda Imtiaz¹, Naghmana Iftikhar², Hina Sattar³, Noshina Riaz⁴

Abstract

Summary: Cushing's syndrome is a condition of excess cortisol activity arising from internal overproduction or external administration of glucocorticoids, leading to a wide range of clinical features.¹ Cushing's syndrome in infancy is rare and most often iatrogenic. We report a case of an infant who presented with rapid-onset weight gain and hypertrichosis due to chronic application of an over-the-counter steroid-containing ointment for diaper rash. Recognition of exogenous steroid exposure allowed for timely withdrawal and clinical recovery. This case highlights the importance of medication history and regulation of topical corticosteroid sales in resource-limited developing countries.

Keywords: Cushing Syndrome; Glucocorticoids; Administration, Topical; Infant; Iatrogenic Disease; Adrenal Insufficiency.

Introduction

Infants are particularly vulnerable to developing topical corticosteroid-induced iatrogenic Cushing's syndrome because of their thin, highly permeable skin, relatively greater surface area-to-body weight ratio, and frequent occurrence of dermatologic conditions requiring topical therapy. Most reported iatrogenic cases have involved the use of potent preparations, such as clobetasol or betamethasone, for diaper dermatitis.

The clinical manifestations of iatrogenic Cushing's syndrome are similar to those of endogenous Cushing's syndrome. Characteristic findings include moon-shaped facies, abnormal weight gain, truncal adiposity with supraclavicular and subscapular fat accumulation, excessive hair growth, facial plethora, and skin fragility with telangiectatic changes. Complications related to mineralocorticoid or androgenic activity, such as hypertension, acne, and hirsutism, are less frequently encountered than those of endogenous diseases. Abrupt cessation of exogenous steroid exposure may precipitate acute adrenal insufficiency, which can be life-threatening if it goes unrecognized.

Case Presentation

A previously healthy 9-month-old female infant presented to the Paediatric Medicine outpatient clinic with progressive weight gain and facial puffiness for two months. The parents initially perceived the chubby cheeks as healthy growth but became concerned as the weight continued to rise rapidly. They also noticed excessive hair growth on the forehead, nape of the neck, periauricular area, and trunk. There was no history of irritability, vomiting, seizures, polyuria, polydipsia, abdominal pain, cough, dyspnoea, oedema, or jaundice. There was no history of systemic steroid use. Birth History was unremarkable, with a birth weight of 3 kg and an uneventful perinatal course. The patient's medical History was also insignificant. The developmental milestones were appropriate for their age. The patient was fully immunised. She was breastfed until 6 months of age, and from 6 months onwards, she was on complementary feeds. On further enquiry, the parents revealed the daily use of a commercially available nappy rash ointment for 4 months. The cream was later identified to contain a potent fluorinated corticosteroid, clobetasol.

On examination, the child was active, playful, and cushingoid, with rounded plethoric facies, central obesity, and hypertrichosis. She was vitally stable with a heart rate of 120/min, respiratory rate 34/min, and BP of 80/40 mmHg (normal for age). There were no striae, acne, skin thinning, oedema, hypertension, hepatosplenomegaly, or features of virilisation.

Contributions:

VI NI - Conception, Design
NI HS NR - Acquisition, Analysis, Interpretation
VI NI - Drafting
HS NR - Critical Review

All authors approved the final version to be published & agreed to be accountable for all aspects of the work.

Conflicts of Interest: None

Financial Support: None to report

Potential Competing Interests:

None to report

Institutional Review Board

Approval

Holy Family Hospital, Rawalpindi

Review began 21/10/2025

Review ended 23/01/2026

Published 31/01/2026

© Copyright 2026

Imtiaz et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY-SA 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



How to cite this article: Imtiaz V, Iftikhar N, Sattar H, Riaz N. Topical Steroid Misuse: An Unrecognized Cause Of Iatrogenic Cushing Syndrome In Infancy. JRM. 2026 Feb. 14;1(1).

<https://doi.org/10.37939/jrmc.v1i1.3064>



Differential Diagnosis

- Iatrogenic Cushing's syndrome due to exogenous steroid use
- Adrenocortical tumour (adenoma/carcinoma)
- Pituitary-dependent Cushing's disease
- Ectopic ACTH production
- Obesity/metabolic syndrome

Investigations

Morning serum ACTH was quite suppressed, that is, 1.5 pg/mL (reference 7.2–63.3 pg/mL), while random blood sugar, serum electrolytes, thyroid profile, HbA1c, and lipid profile were within normal limits. No imaging was performed as adrenal pathology was unlikely, given the suppressed ACTH levels and clear drug history.

Treatment

The offending topical steroid ointment was immediately discontinued. Oral Hydrocortisone was started in physiological doses, that is, 10 mg/m²/day in three divided doses, as patients can experience adrenal crises if steroids are withdrawn immediately. Parents were counselled to avoid unsupervised steroid use. Supportive skin care for diaper dermatitis was also advised.

Outcome And Follow-Up

At the 4-week follow-up, the patient showed gradual regression of cushingoid facies, reduction in hypertrichosis, and reduction in weight. The vital signs and developmental progress remained normal. Long-term follow-up was planned to monitor hypothalamic–pituitary–adrenal axis recovery and growth parameters.



Discussion

In developing regions, the unsupervised or inappropriate use of pharmacological agents remains prevalent because of inadequate regulation of over-the-counter drug sales and the use of potent corticosteroid-containing ointments for diaper rash. Management centres on the gradual withdrawal of the exogenous steroid, treatment of the primary dermatologic condition, and monitoring for adrenal insufficiency during recovery. Public health measures, including the regulation of over-the-counter steroid sales and parental education, are crucial for preventing such cases. Paediatricians should maintain a high index of suspicion for steroid-induced Cushing's syndrome in infants presenting with cushingoid appearance, particularly if topical preparations have been used in the diaper area. Caregivers should always be questioned about the use of creams or ointments applied to the skin.³

Learning Points:

- Always obtain a detailed drug history, including topical agents, in any child with Cushingoid features.
- Topical steroids under occlusion in infants can lead to systemic Cushing's syndrome.
- Early recognition and gradual withdrawal of the offending agent usually result in the reversal of clinical features without long-term sequelae.
- There is an urgent need for stricter regulations on the use of potent topical corticosteroids in resource-limited settings.

Author Information

1. Senior Registrar, Holy Family Hospital, Rawalpindi 2,4. Postgraduate Trainee, Holy Family Hospital, Rawalpindi 3. Associate Professor/Head of Department, Holy Family Hospital, Rawalpindi 4. Vice Chancellor, Rawalpindi Medical University, Rawalpindi.

Corresponding author: Dr. Verda Imtiaz, verdaimtiaz@hotmail.com

References

1. Kuikel S, Aryal S, Basnyat RS, Rimal S. Cushing syndrome in a pediatric patient with topical steroid overuse. *Case Rep Endocrinol.* 2022;2022:8487737. <https://doi.org/10.1155/2022/8487737>

Clinical Case Reports

2. Ahmed SM, Ahmed SF, Othman S, Abdulla BA, Mohammed SH, Salih AM, Kakamad FH. Topical corticosteroid-induced iatrogenic Cushing syndrome in an infant: a case report with literature review. *Ann Med Surg (Lond)*. 2021;71:102978. <https://doi.org/10.1016/j.amsu.2021.102978>
3. Kilci F, Sankaya E. Pediatric iatrogenic Cushing's syndrome: a series of seven cases induced by topical corticosteroid use. *J Pediatr Endocrinol Metab*. 2025;38(8):863–867. <https://doi.org/10.1515/jpem-2025-0032>
4. Matejek N, Hoos J, Holterhus PM, Menke LA, Lüdecke HJ, Bettendorf M, et al. Topical glucocorticoid application causing iatrogenic Cushing's syndrome followed by secondary adrenal insufficiency in infants: two case reports. *J Med Case Rep*. 2022;16:455. <https://doi.org/10.1186/s13256-022-03659-2>