

## Endocrine Dysregulation and Functional Ovarian Cyst: A Case from Adolescent Gynecology

Tabinda Khalid<sup>1</sup>, Rubaba Abid Naqvi<sup>2</sup>, Aqsa Ikram Ul Haq<sup>3</sup>, Shama Bashir<sup>4</sup>

### Abstract

**Summary:** Functional ovarian cysts are common in reproductive-age women, but their occurrence secondary to hypothyroidism is rare. We report a case of a 21-year-old female with poorly controlled hypothyroidism who presented with bilateral multiloculated ovarian cysts mimicking neoplastic pathology. Emergency laparotomy was performed due to worsening symptoms, and bilateral cystectomy with ovarian conservation was achieved. Histopathology confirmed corpus luteal cysts. Postoperative thyroid hormone replacement led to normalization of menstrual cycles and resolution of cysts. This case highlights the importance of thyroid evaluation in young women with ovarian masses to avoid unnecessary surgery and preserve fertility.

**Keywords:** Ovarian Cysts, Corpus Luteum, Hypothyroidism, Thyroid Hormones/therapeutic use, Adolescent, Young Adult, Gynecology, Fertility Preservation.

### Introduction

Ovarian cysts are fluid-filled sacs that may arise from normal ovulatory processes or pathological conditions. Ovarian cysts are a common gynecologic finding in adolescents, yet their evaluation and management require careful consideration due to age-specific risks and outcomes.<sup>1</sup> Updated guidelines emphasize structured approaches for diagnosis and treatment in children and adolescents, ensuring both safety and fertility preservation.<sup>2</sup> Recent retrospective studies further highlight the clinical spectrum of adolescent ovarian cysts and the importance of individualized management strategies.<sup>3</sup> Endocrine disorders like hypothyroidism can disrupt the hypothalamic–pituitary–ovarian axis.<sup>4</sup>

It may affect ovarian function through complex endocrine and immune pathways.<sup>5</sup> Recognition of this link is crucial, as thyroid hormone replacement can reverse changes and restore reproductive function.<sup>6</sup>

### Case Presentation

A 21-year-old unmarried female, a postgraduate student from Rawalpindi, presented to the Gynecology outpatient department of Rawalpindi Teaching Hospital on 11 April 2025 with complaints of lower abdominal pain ongoing for two years, with acute worsening over the preceding week, along with menstrual irregularities. She had a three-year history of hypothyroidism but was non-compliant with levothyroxine therapy. Her menstrual cycles were previously regular (5/28); however, for the past two years, she had become oligomenorrheic, with scanty flow every 40–60 days. She reported progressive weight gain but denied cold intolerance, constipation, acne, hirsutism, galactorrhea, or urinary and bowel disturbances. On examination, she was well-oriented with a BMI of 29.8 kg/m<sup>2</sup>, pulse rate 106/min, and blood pressure 100/60 mmHg. The abdomen was protuberant with a palpable, firm, non-tender, irregular pelvic mass measuring approximately 20 cm, with restricted mobility.

#### Investigations:

- Hemoglobin: 10.1 g/dL
- TSH: 41 mIU/L (↑), Free T4: low
- CA-125: 18.4 IU/mL
- Ultrasound: Bilateral multilocular adnexal cysts- right cyst (16x11x4) and left cyst (10 × 11 × 5 cm)—with normal uterus.
- CT scan: Confirmed complex cysts with mild ascites
- Tumor markers: CA-125 18.4 IU/ml, β-hCG < 1 IU/ml, AFP 5 IU/ml.

#### Treatment

We planned a laparoscopy; however, due to worsening pain, an emergency laparotomy was performed. Bilateral multilocular cysts with hemorrhagic areas were found. Bilateral cystectomy with ovarian conservation was completed. Histopathology confirmed corpus luteal cysts. Levothyroxine 100 µg/day was initiated postoperatively.

#### Outcome and Follow-Up

At 3-month follow-up:

- Menstrual cycles normalized
- Ultrasound showed normal ovaries
- TSH decreased to 4 mIU/L

#### Contributions:

TK - Conception, Design  
TK - Acquisition, Analysis, Interpretation  
TK RAN AIUH SB - Drafting  
TK RAN AIUH SB - Critical Review

All authors approved the final version to be published & agreed to be accountable for all aspects of the work.

#### Conflicts of Interest: None

**Financial Support:** None to report

#### Potential Competing Interests:

None to report

#### Institutional Review Board

##### Approval

25/RTH, RWP

02-01-2026

Rawalpindi Teaching Hospital

Review began 12/12/2025

Review ended 01/01/2026

Published 31/01/2026

© Copyright 2026

Khalid et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY-SA 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**How to cite this article:** Khalid T, Naqvi RA, Haq AIU, Bashir S. Endocrine Dysregulation and Functional Ovarian Cyst: A Case from Adolescent Gynecology. JRMC. 2026 Feb. 14;1(1).

<https://doi.org/10.37939/jrhc.v1i1.3197>



Figure 1A: (ultrasound image showing multiloculated ovarian cysts)



Figure 1B: (C-T Scan showing bilateral complex cysts with ascites)

## Discussion

Functional ovarian cysts arise from disturbances in normal follicular development and are most common in reproductive-age women. The corpus luteal cysts observed in this patient are typically transient structures that may enlarge due to hormonal dysregulation. Primary hypothyroidism has been implicated in their pathogenesis through complex interactions between the hypothalamic–pituitary–ovarian axis. In hypothyroidism, several mechanisms can lead to the development of large ovarian cysts.




Figure 1 C - (Preserved ovaries after cyst removal)

Elevated thyroid-stimulating hormone (TSH) can cross-react with follicle-stimulating hormone (FSH) receptors on ovarian tissue, promoting excessive follicular growth. Increased thyrotropin-releasing hormone (TRH) causes hyperprolactinemia, which disrupts gonadotropin-releasing hormone (GnRH) pulsatility, leading to anovulation and follicular persistence. Additionally, altered gonadotropin ratios with relatively low luteinizing hormone (LH) and higher FSH levels prevent normal luteinization, resulting in cyst formation. Chronic hypothyroidism may also increase ovarian sensitivity to gonadotropins and insulin-like growth factors, further stimulating cyst development. Lastly, thyroid hormone deficiency directly impairs ovarian follicular maturation and angiogenesis, contributing to the persistence of large functional cysts.

Hypothyroidism presenting with ovarian cysts is an uncommon but clinically important condition. Large cysts may mimic neoplastic disease, creating diagnostic uncertainty and the risk of unnecessary surgical intervention. Similar to our patient, huge bilateral ovarian cysts associated with hypothyroidism have been reported, highlighting the potential for misinterpretation as ovarian malignancy.<sup>7</sup>

Other reports describe bilateral ovarian masses due to primary hypothyroidism, further emphasizing the spectrum of presentations and the importance of considering endocrine causes in the differential diagnosis.<sup>8</sup> Rare complications such as pituitary adenoma and chronic adnexal torsion have also been documented in hypothyroid patients, underscoring the need for timely recognition and multidisciplinary management.<sup>9</sup>

Additionally, giant ovarian cysts have been described following thyroidectomy, reinforcing the link between thyroid dysfunction and ovarian pathology.<sup>10</sup> Collectively, these cases demonstrate that ovarian cysts secondary to hypothyroidism can regress with appropriate thyroid hormone replacement, thereby avoiding unnecessary surgical procedures. Our case adds to this growing body of evidence, supporting routine thyroid evaluation in adolescents and young women presenting with ovarian cysts.

Collectively, these cases demonstrate that ovarian cysts secondary to hypothyroidism can regress with appropriate thyroid hormone replacement, thereby avoiding unnecessary surgical procedures. Our case adds to this growing body of evidence, supporting routine thyroid evaluation in adolescents and young women presenting with ovarian cysts. Thus, a multidisciplinary team approach involving gynecologists, endocrinologists, and radiologists is essential for accurate diagnosis, timely intervention, and fertility preservation, as functional cysts secondary to hypothyroidism are reversible with appropriate hormonal supplementation.

## Conclusion

Functional ovarian cysts secondary to hypothyroidism, though uncommon, should be considered in reproductive-age females presenting with bilateral ovarian masses. Early recognition and thyroid hormone replacement can lead to complete regression of cysts and normalization of reproductive function, thereby avoiding unnecessary surgical procedures and preserving fertility.

### Learning points

- Thyroid function testing should be routine in young women presenting with bilateral ovarian cysts.
- Functional cysts secondary to hypothyroidism may mimic malignancy but regress with thyroid hormone therapy.
- Multidisciplinary management involving gynecologists, endocrinologists, and radiologists is essential to prevent unnecessary surgery and preserve fertility

## Author Information

1. Senior Registrar, Gynae/Obs, RTH/DHQ Hospital, Rawalpindi 2. Professor, Gynae/Obs, RTH/DHQ Hospital, Rawalpindi 3,4. Assistant Professor, Gynae/Obs, RTH/DHQ, Rawalpindi.

**Corresponding author:** Dr. Tabinda Khalid, tabindakhalid@gmail.com

## References

1. Bašković M, Habek D, Zaninović L, Milas I, Pogorelić Z. The evaluation, diagnosis, and management of ovarian cysts, masses, and their complications in fetuses, infants, children, and adolescents. *Healthcare (Basel)*. 2025;13(7):775. <https://doi.org/10.3390/healthcare13070775>
2. Ritchie J, O'Mahony F, Garden A. Guideline for the management of ovarian cysts in children and adolescents. *Br Soc Paediatr Adolesc Gynecol Guideline*. 2024.
3. Retnowati S, Wahyuni A, Widiartara IN, Nisrina H. Clinical characteristics and management of adolescent ovarian cysts: a retrospective study. *Indones J Obstet Gynecol*. 2025;13(2):2169. <https://doi.org/10.32771/inajog.v13i2.2169>
4. Tan J, Yang YY, Yin DY, Xin Q, Cai GX. Research on the impact of thyroid disorders on reproductive function: a narrative review. *J Clin Med Res*. 2025 Aug;17(8):409-422. <https://doi.org/10.14740/jocmr6315>
5. Popa EC, Maghiar L, Maghiar TA, Brihan I, Georgescu LM, Toderas BA, Sachelarie L, Huniadi A. Hashimoto's thyroiditis and female fertility: endocrine, immune, and microbiota perspectives in assisted reproduction—a narrative review. *Biomedicines*. 2023;13(6):1495. <https://doi.org/10.3390/biomedicines13061495>
6. Zhang H, Qiu H, Liu Z, Wu Y, Liu W, Huang C. Subclinical/overt hypothyroidism may be associated with diminished ovarian reserve in infertile women independent of thyroid autoimmunity. *Front Endocrinol*. 2024 Dec;15:1477665.
7. Jain D, Jain S. Huge bilateral ovarian cysts with concurrent hypothyroidism: a case report. *Cureus*. 2024 Apr 23;16(4):e58837. <https://doi.org/10.7759/cureus.58837>
8. Atak Z, Oral B, Güneş E, Ocakoğlu SR, Bayram F. A case of primary hypothyroidism complicated with pituitary pseudo-macroadenoma and chronic adnexal torsion. *Cyprus J Med Sci*. 2024;9(6):450-453. <https://doi.org/10.4274/cjms.2024.2024-58>
9. Kodra B. Resolution of ovarian cysts in reproductive-age women with hypothyroidism: a case series and review of the literature. *Int J Basic Life Sci*. 2025;13(2). <https://doi.org/10.22034/ijbls.2025.223570>
10. Monteiro JM, Campos M, Estevinho N, Vasconcelos-Castro S. Giant ovarian cyst after total thyroidectomy: a case report. *Surg Gastroenterol Oncol*. 2024 Jul;29(2 Suppl). <https://doi.org/10.21614/sgo-685>